

EMERGENCY RECORD	Date		
Student	Home Phone		
Grade Gender	Birth Date		
Child lives with: Both Parents Mother	Father Other		
Please use area code and identify phone number types: W = Work	C = Cell E = Evening		
Parent or Guardian 1	Relationship to Student		
Phone 1 Phone 2 Phone 2	Phone 3 Child Lives With		
Email	Step-parent		
Address (if different than above)			
Parent or Guardian 2	Relationship to Student		
Phone 1 Phone 2	Phone 3 Child Lives With		
Email	Step-parent		
Address (if different than above)			
Immunizations within the last year for returning students (Type, mo	/day/yr)		
Chronic condition(s)			
Allergies			
Medications			
IN CASE OF EMERGENCY			
Contact 1	Relationship to Student		
Phone 1 Phone 2	Phone 3		
Contact 2	Relationship to Student		
Phone 1 Phone 2	Phone 3		
In case of illness or mild accidents, our procedure will be to cordesignated on this emergency record. In case of a severe emer an emergency care center. We will attempt to call you as soon	gency, an ambulance will be called to take your child to		



POTASSIUM IODIDE PARENT/GUARDIAN INFORMED CONSENT FORM

Reason for Taking Potassium Iodide

In the case of an accident at a nuclear power plant or what is known as a radiological emergency, radioactive iodine will be released into the air. This material may be inhaled or ingested and enter the thyroid gland where it can cause cancer and/or disease. Children and infants are the most vulnerable to this occurrence. When taken by pill or liquid, potassium iodide (KI) floods the thyroid with non-radioactive iodine and prevents the thyroid from absorbing radioactive material. Potassium iodide needs to be given before or shortly after exposure to radiation. Potassium Iodide works only to prevent the thyroid from absorbing radioactive iodine.

Potential Side Effects of Potassium Iodide

It is possible to experience any or all of the following side effects when taking Potassium Iodide:

- Allergic reaction
- Upset Stomach
- Rash

Risks of Taking Potassium Iodide

Taking Potassium Iodide is safe for most people. Potassium Iodide should not be taken if someone:

- Is allergic to iodine
- Has Graves Disease or any other thyroid illness
- Takes thyroid medication
- Has Dermatitis Herpetiformis or Hypocomplementemic Vasculitis

Administration of Potassium Iodide

The recommended dose of Potassium Iodide is 1 tablet (65 mg) for children 3 years and up who weigh less than 150 pounds and 2 tablets (130 mg) for children over 12 who weigh at least 150 pounds. The recommended dose of Potassium Iodide will only be given:

- In case of a radiological emergency
- Directed by public health officials
- If a parent/guardian signs a consent form for a child

I consent to have Holy Sp	oirit Academy admir	nister Potassium Iodide (KI)	to my child.
	Yes	No	
I have read and understand the in	nformation on this p	page regarding Potassium Io	dide administration.
Parent or Guardian Signature		D	ate